



## DEMOGRAPHICS

Total population (000)	20,714	(2010)
Total under-five population (000)	3,305	(2010)
Births (000)	732	(2010)
Birth registration (%)	80	(2008-2009)
Total under-five deaths (000)	44	(2010)
Neonatal deaths: % of all under-5 deaths	37	(2010)
Neonatal mortality rate (per 1000 live births)	22	(2010)
Infant mortality rate (per 1000 live births)	43	(2010)
Stillbirth rate (per 1000 total births)	21	(2009)
Total maternal deaths	1,800	(2010)
Lifetime risk of maternal death (1 in N)	81	(2010)
Total fertility rate (per woman)	4.7	(2010)
Adolescent birth rate (per 1000 women)	147	(2007)

### Under-five mortality rate




### Maternal mortality ratio



Note: MDG target calculated by Countdown to 2015


## MATERNAL AND NEWBORN HEALTH

### Coverage along the continuum of care




### Skilled attendant at delivery

Percent live births attended by skilled health personnel



### Prevention of mother-to-child transmission of HIV


Eligible HIV+ pregnant women receiving ART for their own health (% of total ARVs) 6 (2010)



## EQUITY

### Socioeconomic inequities in coverage

Household wealth quintile: ● Poorest 20% ● Richest 20%




Coverage levels are shown for the poorest 20% (red circles) and the richest 20% (orange circles). The longer the line between the two groups, the greater the inequality. These estimates may differ from other charts due to differences in data sources.

## CHILD HEALTH


### Immunization

■ Percent of children immunized against measles  
■ Percent of children immunized with 3 doses DTP  
■ Percent of children immunized with 3 doses Hib



### Pneumonia treatment

■ Percent children <5 years with suspected pneumonia taken to appropriate health provider  
■ Percent children <5 years with suspected pneumonia receiving antibiotics




## NUTRITION

Wasting prevalence (moderate and severe, %)	15 (2003-2004)	Early initiation of breastfeeding (within 1 hr of birth, %)	72 (2008-2009)
Low birthweight incidence (moderate and severe, %)	16 (2008-2009)	Introduction of solid, semi-solid/soft foods (%)	86 (2008-2009)
		Vitamin A supplementation (two dose coverage, %)	95 (2010)

### Underweight and stunting prevalence


■ Percent children <5 years who are underweight  
■ Percent children <5 years who are stunted



Note: Based on 2006 WHO reference population


### Exclusive breastfeeding

Percent infants <6 months exclusively breastfed




## DEMOGRAPHICS

### Causes of under-five deaths, 2010



\*Intrapartum-related events \*\*Sepsis/meningitis/tetanus

### Causes of maternal deaths, 1997-2007




## POLICIES

Maternity protection in accordance with Convention 183	No
Specific notification of maternal deaths	Yes
Midwifery personnel authorized to administer core set of life saving interventions	Yes
International Code of Marketing of Breastmilk Substitutes	Yes
Postnatal home visits in first week of life	Yes
Community treatment of pneumonia with antibiotics	Yes
Low osmolarity ORS and zinc for management of diarrhoea	Partial
Rotavirus vaccine	-
Pneumococcal vaccine	Partial

## MATERNAL AND NEWBORN HEALTH

### Antenatal care

Percent of women aged 15-49 years attended at least once by a skilled health provider during pregnancy



Demand for family planning satisfied (%)	60	(2008-2009)
Antenatal care (4 or more visits, %)	49	(2008-2009)
Malaria during pregnancy - intermittent preventive treatment (%)	6	(2008-2009)
C-section rate (total, urban, rural; %)	2, 6, 1	(2008-2009)
(Minimum target is 5% and maximum target is 15%)		
Neonatal tetanus vaccine (%)	76	(2010)
Postnatal visit for baby (within 2 days for all births, %)	-	-
Postnatal visit for mother (within 2 days for all births, %)	46	(2008-2009)
Women with low body mass index (<18.5 kg/m <sup>2</sup> , %)	28	(2008-2009)


## SYSTEMS AND FINANCING

Costed national implementation plan(s) for maternal, newborn and child health available	Yes
Density of doctors, nurses and midwives (per 10,000 population)	4.8 (2007)
National availability of emergency obstetric care services (% of recommended minimum)	11 (2010)
Per capita total expenditure on health (Int\$)	65 (2010)
General government expenditure on health as % of total government expenditure (%)	15 (2010)
Out-of-pocket expenditure as % of total expenditure on health (%)	27 (2010)
Official development assistance to child health per child (US\$)	10 (2009)
Official development assistance to maternal and neonatal health per live birth (US\$)	17 (2009)

## CHILD HEALTH

### Diarrhoeal disease treatment


- Percent of children <5 years with diarrhoea receiving oral rehydration therapy/increased fluids with continued feeding
- Children <5 years with diarrhoea treated with ORS



### Malaria prevention and treatment

Percent of children receiving first line treatment among those receiving any antimalarial

- Percent of children <5 years sleeping under ITNs




## WATER AND SANITATION

### Improved drinking water coverage

Percent of population by type of drinking water source, 1990-2010

- Piped on premises
- Other improved
- Unimproved
- Surface water




Source: WHO/UNICEF JMP 2012

### Improved sanitation coverage

Percent of population by type of sanitation facility, 1990-2010

- Improved facilities
- Shared facilities
- Unimproved facilities
- Open defecation



Source: WHO/UNICEF JMP 2012